References

Akinbowale, O. L., Peng, H., Barton, M. D., 2006. Antimicrobial resistance in bacteria isolated from aquaculture sources in Australia. J. Appl. Microbiol. 100, 1103–1113.

Arnow, L. E., 1937. Colorimetric determination of the components of 3,4-dihydroxyphenylalanine –tyrosine mixtures. J Biol Chem 118, 531 –537.

Atkin, C. L., Neilands, J. B., Phaff, H. J., 1970. Rhodotorulic acid from species of Leucosporidium, Rhodosporidium, Rhodotorula, Sporidiobolus, and Sporobolomyces, and a new alaninecontaining ferrichrome from Cryptococcus melibiosum. J Bacteriol 103, 722 –733.

Atlas, R. M., Bartha, E., 1997. Microbial ecology fundamentals and application. Benjamin Cummings Science Publishing, Menlo Park, CA.

Das, B. K., Samal, S. K., Samantaray, B. R., Sethi, S., Pattnaik, P., Mishra, B. K., 2006. Antagonistic activity of cellular components of Pseudomonas species against Aeromonas hydrophila. Aquaculture. 253, 17–24.

Diez-Gonzalez, F., 2007. Use of bacteriocin in livestock. In: Riley M. A., Gillor O. (Eds.), Research and applications in bacteriocins. Horizon Bioscience, Norfolk, pp. 117–129.

Dong, Y. H., Zhang, L. H., 2005. Quorum sensing and quorum-quenching enzymes. J. Microbiol. 43, 101-109.

Gatesoupe, F. J., 1999. The use of probiotics in aquaculture. Aquaculture 180, 147-165.

Havenaar, R., Ten Brink, B., Huis Int Veld, J. H. J., 1992. Selection of strains for probiotic use. In: Fuller. R. (Ed.). Probiotics the Scientific Basis. Chapman and Hall, London, pp. 209—229.

Lembke, C., Podbielski, A., Hidalgo-Grass, C., Jonas, L., Hanski, E., Kreikemeyer, B., 2006. Characterization of biofilm formation by clinically relevant serotypes of group A streptococci. Appl. Environ. Microbiol. 72, 2864-2875.

Maniatis, T., Fritsch, E. F., Sambrook, J., 1982. Molecular Cloning: A Laboratory Manual. Cold Spring Harbour Laboratory. Cold Spring Harbour, New York.

Moriarty, D. J. W., 1998. Control of luminous Vibrio species in penaeid aquaculture ponds. Aquaculture 164, 351-358.

Moriarty, D. J. W., 1999. Disease control in shrimp aquaculture with probiotic bacteria. In: Microbial Biosystems: New Frontiers. In: Bell, C. R., Brylinsky, M., Johnson-Green, P. (Eds.), Proceeding of the 8th International Symposium on Microbial Ecology, Atlantic Canada Society for Microbial Ecology, Halifax, Canada, pp. 237-243.

Neilands, J. B., 1981. Microbial iron transport compounds (siderophores) as chelating agents. In: Development of iron chelators for clinical use, Martell, A. E., Anderson. W. J. and Badman, D.G. (Eds.), Elsevier, North Holland, Amsterdam, pp.13-31.

Ouwehand, A. C., Kirjavainen, P. V., Gronlund, M. M., Isolauri, E., Salminen, S., 1999. Adhesion of probiotic micro-organisms to intestinal mucus. Int. Dairy J., 9, 623-630.

Pan, X., Wu, T., Zhang, L., Song, Z., Tang, H., Zhao, Z., 2008. In vitro evaluation on adherence and antimicrobial properties of a candidate probiotic Clostridium butyricum CB2 for farmed fish. J. Appl. Microbiol. 105, 1623– 1629.

Payne, S. M., 1994. Detection, isolation and characterization of siderophores. Method Enzymol. 235, 329–344.

Rode, T. M., Langsrud, S., Holck, A., Møretrø, T., 2007. Different patterns of biofilm formation in Staphylococcus aureus under food-related stress conditions. Int. J. Food Microbiol. 30, 372-83.

Schwarz, S., Kehrenberg, C., Walsh, T. R., 2001. Use of antimicrobial agents in veterinary medicine and food animal production. Int. J. Antimicrob. Ag. 17, 431–437.

Schwyn, B., and Neilands, J. B., 1987. Universal chemical assay for the detection and determination of siderophores. Anal Biochem 160, 47 –56.

Shenker, M., Oliver, I., Helmann, M. , Hadar, Y., Chen. Y., 1992. Utilization by tomatoes of iron mediated by a siderophore produced by Rhizopus arrhizus. Journal of Plant Nutrition, 10, 2173-82.

Snow, G. A., 1954. Mycobactin. A growth factor for Mycobacterium johnei. Part II. Degradation and identification of fragments. J. Chem. Soc. 2588-2596.

Stewart, J. E. 1965. Far infrared spectroscopy. In S. K. Freeman (ed.), Interpretive spectroscopy. Reinhold Publishing Corp., New York, pp. 131-169.

Thenmozhi, R., Nithyanand, P., Rathna, J., Pandian, S. K., 2009. Antibiofilm activity of coral associated bacteria against different clinical M serotypes of Streptococcus pyogenes. FEMS Immunol. Med. Microbiol. 57, 284-294.

Trevors, J. T., 1986. Plasmid curing in bacteria. FEMS Microbiology Letters, 32, 149–157.

Verschuere, L., Rombaut, G., Sorgeloos, P., Verstraete, W., 2000. Probiotic bacteria as biological control agents in aquaculture. Microbiol. Mol. Biol. R. 64, 655-671.

Vesterlund, S., Paltta, J., Karp, M., Ouwehand, A. C., 2005. Measurement of bacterial adhesion - in vitro evaluation of different methods. J. Int. Microb. Meth. 60, 225-233.

Vine, N. G., Leutes, W. D., Kaiser, H., Daya, S., Baxter, J., and Hecht, T., 2004. Competition for attachment of aquaculture candidate probiotic and pathogenic bacteria on fish intestinal mucus. J. Fish Dis. 27, 319-326.

National and International conferences

1. **Aparna Balakrishna and T R Keerthi,** "Characterization of siderophore produced by the gut microflora of *Poecilia reticulata* for potential use as probiotic in aquaculture", World Congress on Biotechnology, Mar 21-23, 2011, Hyderabad, India.

2. **Aparna Balakrishna and T R Keerthi, "**Screening of Potential Aquatic Probiotic from Major Microflora of Guppy (*Poecilia reticulata*)", Asian Congress on Biotechnology 2011, May 11-15, 2011, Shanghai, China.

Projects undertaken

1. Project entitled "Production Purification and characterization of Antagonistic principles of potential Aquaculture Probiotics", funded by University Grants Commission (UGC), New Delhi, India under grant UGC-F.No.37-1/2009 (KER) (SR).

Papers submitted

1. Aparna Balakrishna and T R Keerthi, "Screening of Potential Aquatic Probiotic from Major Microflora of Guppy (*Poecilia reticulata*)", submitted to *Front. Chem. Sci. Eng.*